Development and Evaluation of Antifungal Nasal Herbal formulation.

A Synopsis submitted to

IIMT University in partial fulfillment of the requirements for the

Degree of

DOCTOR OF PHILOSOPHY

In

Pharmaceutical Sciences

By

Archana Gautam

Under the Supervision of

Dr. ANUJ MITTAL
(Associate Professor)
Department of Pharmacy

IIMT COLLEGE OF MEDICAL SCIENCES,
IIMT UNIVERSITY, MEERUT, UTTAR PRADESH

DECLARATION

I hereby declare that this synopsis report submission is my own work and that, to the best of my knowledge and belief, it contains no material previously published or written by another person nor material which to a substantial extent has been accepted for the award of any other degree or diploma of the university or other institute of higher learning, except where due acknowledgment has been made in the text.

Archana Gautam

Roll No. 2165004004

Dr. Divya Pathak
(Associate Professor)
IIMT College of Medical Sciences,

(Supervisor)

Date:
Place: Meerut

TABLE OF CONTENTS

S. No.	TITLE	PAGE NO.
1	INTRODUCTION	1-10
2	LITERATURE REVIEW	11-12
3	AIM & OBJECTIVES	13
4	PLAN OF WORK	14-15
5	EXPECTED OUTCOMES	16
6	REFERENCES	17-18

INTRODUCTION

Inflammation of the nasal mucous membrane and frequently the conjunctiva mucous membrane is known as allergic rhinitis (AR). It develops as a result of exposure to inhaled allergenic substances and causes an immunoglobulin-mediated response (IgE**) (Dipiro et al., 2002; Shargel et al., 2001)**. Four main symptoms, or major primary symptoms, define AR. Some signs include sneezing, wet rhinorrhoea, nasal blockage, and itching **(Min,2010)**. Postnasal drip, itchy eyes, ears, nose, and palate are other common symptoms of AR in addition to the primary ones (Kemp, 2009). Around two out of every ten persons worldwide are affected by AR, which occurs regularly (Weinberg, 1993). With an incidence of 10-25% and a prevalence that is rising and affects 5–50% of the world's population, AR is extremely prevalent worldwide **(Braido et al., 2008; Lee et al., 2008; Wilson et al., 2002).** Although AR does not pose a life-threatening hazard, it has a major negative impact on quality of life **(Nelson 2007; Wilson et al., 2002).**

1.1 Background

The therapeutic objective for patients with AR is to avoid or reduce symptoms utilising a course of treatment with little to no side effects and at a fair cost, allowing patients to lead normal lives **(Dipiro et al.,2002).** Antihistamines, decongestants, antihistamine decongestant combos, anticholinergics, corticostreoids, ipratropium bromide, leukotriens modifiers, mast cell stabilisers, and immunotherapy are all included in conventional treatment. Respiratory allergies are one of the most common conditions for which homoeopathic treatment is sought after. It has been suggested that the herbal treatments Alliemcepa, Euphrasiaofficinalis, Luffaoppercuata, and Sabadillaofficinaraum included in Otirin Nasal Spray may be useful in treating allergic rhinitis. A homoeopathic complex product called otirin nasal spray is recommended for AR. The effectiveness of this product in treating this ailment has not been specifically studied. thus, our research is primarily concentrated on the creation of a poly-herbal formulation employing Moringa oliefera and Embelia ribes extracts.

1.2 Allergic rhinitis

Sneezing, nasal congestion, a runny or stuffy nose, and inflammation of the nasal mucosal membranes are all symptoms of allergic rhinitis (AR). According to a review of the literature, mediators that come into contact with the nose and eye linings—such as pollen, mould, dust, mites, and animal allergens—cause inflammation of the nasal mucosal membranes. Since these mediators are thought to play a role in the pathogenesis of AR, it is believed that these targets can be further investigated in order to look into other formulations that might be utilised to treat the disorders' symptoms. According to a thorough study of the literature, no poly-herbal nasal preparation using particular plants has ever been created to treat AR. So, the goal of the current study is to create poly-herbal nasal formulations and evaluate their acceptability in mice who have had their ovaries damaged in order to treat AR. As a result of a transient or ongoing allergic reaction to allergens, AR is a type of IgE-mediated inflammatory cell that irritates the airway tissue layer. In addition to nasal sneezing, rubbing, rhinorrhea, tears, congestion, and obstruction, AR is primarily characterised by an area flow of eosinophils. AR also causes overall irritability and is linked to a number of co-morbid conditions, including respiratory disorders, nasal polypsis, rhino sinusitis, humour otitis, and sleep issues. Mast cell-mediated allergic reactions make up the majority of the inflammatory response in the nasal tissue layer, while a delinquent part of the response is defined by the inflammatory recruitment of secondary effector cells, such as eosinophils, basophils, and T cells.

1.3 Allergens

An allergen is an antigenic material that causes an allergic reaction or hypersensitive reaction (Martini 2004; Stedman, 2008). Exposure to seasonal and perennial allergens that are present in both indoor and outdoor environments frequently causes AR. IgE-mediated immunological pathways play a significant role in the development of allergy symptoms by causing the release of mediators (Wang 2005).

1.4 Nasal Anatomy and physiology

The primary air entry point for the respiratory system is the nose, which is part of the respiratory tract. The peripheral organ of smell is also located in the nose. The inferior two thirds of the nasal mucosa make up the respiratory area of the nose, whereas the superior third makes up the olfactory area. The olfactory region houses the peripheral organ of smell (Martini, 2004; Moore and Dalley, 2006). Due to the abundant vascularity of the nasal membrane and the convenience of intranasal administration, the nasal route of administration is a practical and reliable approach for both local and systemic medication delivery. The nasal mucosa is currently being considered as a route of administration to achieve a quicker and healthier amount of medication absorption. Several medications, including proteins and peptides, find the nasal route of administration appealing because to the nasal membrane's rich vascular environment and high drug permeability. Moreover, drug absorption in the nose's olfactory area makes it possible for a pharmaceutical molecule to reach the central nervous system. Another terrible use for vaccinations in terms of effectiveness and patient acceptance is nasal administration. According to custom, mankind has been interested in using the nasal route for medicine delivery from the dawn of time. Nasal therapy, also known as "Nasaya Karma," is a recognised kind of treatment in the Indian Ayurvedic medical system.

REVIEW OF LITERATURE

2.1 General a systemic response to the allergy and underlying inflammation should be treated well in order to address the AR. It is necessary to have a thorough awareness of the pathology, the connection to co-morbid diseases and impacts, as well as the numerous therapy alternatives **(Borish, 2003).** The goal of treatment for AR patients is to avoid or reduce symptoms while employing therapies with few to no side effects and reasonable cost, allowing them to continue a normal way of life. Results of effective treatment include a reduction in ER visits, hospitalisations for asthma attacks, and the requirement for grommets in cases of otitis media with effusion, as well as a marked improvement in the quality of life of AR patients **(Scadding, 2008).**

Antihistamines, decongestants, antihistamine decongestant combos, anticholinergics, corticosteroids, ipratopium bromide, leukotriens modifiers, mast cell stabilisers, herbal extracts used naturally, and immunotherapy were all included in conventional treatment. A specific pharmacotherapy is chosen in accordance with information specific to the patient.

While successful, more effective therapies are still needed, hence in our study we used formulations of poly-herbal extracts with low dose and improved local effects **(Braido et al., 2008).**

(Blaiss, 2004 a) The quality of life for paediatrics may decline significantly, child development may suffer, and there may be an increased risk of developing various illnesses such as bronchial asthma, chronic sinusitis, and other breathing problems.

The Blaiss (2004b)

Mold, pollen, animal dander, dust mites, pollution, and various airborne particles are common causes of allergic rhinitis. These compounds can be found almost everywhere on earth and often do not depend on the season. While some of these allergens are seasonal, others are common all year round.

(Momoe, 2001)

It must be noted that allergic rhinitis is not always communicable and cannot be passed from one person to another, regardless of the circumstance. Every man and woman is susceptible to the virus, which is brought on by a single character predisposition.

(McCabe, 2008)

According to reports, 30–40% of people with allergic rhinitis continue to have symptoms for up to 5 hours after the specific exposure to the allergen that caused the reaction. This occurs in part because the body continues to release chemicals to fight the allergen that has entered the body on a regular basis. 2.2 Allergy rhinitis natural therapy Even while certain prescription therapies are successful in treating AR, many of the commonly prescribed pharmaceuticals can have negative side effects. As a result, people with AR frequently look for more natural treatments**. (Balch et al., 2008).**

(Pratibha et al., 2004)

For the treatment of allergic rhinitis, a brand-new, secure polyherbal compound (Aller-7/NR-A2) has been developed using an entirely original combination of the extracts. In order to treat allergic rhinitis, a novel polyherbal compound (Aller-7/NR-A2) was developed using a special mix of extracts from seven medicinal flowers.

2.3 Complimentary and alternative medicines

(Rao et al., 2009) The anti-inflammatory effects of the aqueous and ethanolic extract of the leaves of M. olifera Lam (Family: Moringaceae) have been investigated in albino rats. The activity responded at its highest level in two extracts. Quercitin is a flavonoid with anti-inflammatory and anti-allergy effects. In preliminary experiments, quercitin has been demonstrated to decrease the release of histamine from mast cells**. (Balch et al., 2008).**

(Anwar et al., 2007)

The most widely farmed tree species in the genus Moringa, which is a member of its own family Moringaceae, is MoringaoleiferaLan (M. oleifera). M. oleifera is a highly prized plant that is found in unique tropics and subtropics areas all over the world.

(Wadhwa 2013)

The moringaoleifera tree is regarded as one of the most useful trees in the world and has a wide range of traditional medicinal, commercial, and dietary uses.

(Makkar and Becker, 1997)

Because they include ascorbic acid, flavonoids, phenolic acids, and carotenoids, the leaves of M. Oleifera are a great source of a natural antioxidant.

(Santos et al., 2011)

The unique elements of the Moringaoleifera tree, including its leaves, roots, seeds, bark, fruits, flora, and immature pods, have been used for treating a variety of human illnesses in traditional medicine, particularly in South Asia. These unique components have antitumor, antipyretic, antiepileptic, anti-inflammatory, antispasmodic, antioxidant, antibacterial, and antifungal properties.

(Tesfay et al., 2011)

The Moringa bushes are naturally found in tropical temperatures in various parts of the world; the extent of their capacity to adapt to cooler climates is tied to metabolic alterations that cause a rise in a variety of natural solutes, including sugars, prolinepolyols, and amino acids.

(Palada and Chang, 2003)

Because it is drought resilient and prefers rainfall levels between 250 and 1500 mm, Moringaoleifera is a fast-growing tree that can tolerate a wide range of environmental conditions. Temperatures between 18.7 to 28.5 oC every 12 months are ideal for growth, while they can go up to 48 oC and cause light frost.

(Fang et al., 2002)

Antioxidants are vital components of the human diet and are important for maintaining health and preventing some diseases. Inhibiting and scavenging free radicals, which can lead to a number of diseases such as most malignancies, heart disease, hypertension, and stroke, are excellent functions of antioxidant sports.

(Siddhuraju and Becker, 2003)

One of the plant species, Moringaoleifera, has a variety of phenolic, flavonoid, and vitamin-based herbal antioxidants.

(Srikanth et al., 2014)

The moringa tree is regarded as one of the most important trees for the nutritional security of rural people; as a result, moringa helps to promote human fitness and fight malnutrition, especially in developing nations.

(Wadhwa, 2013)

Nearly all of the Moringaoleifera tree's components, including the root, bark, leaves, buds, plants, and seeds, have been used in traditional South Asian medicine for a variety of diseases. These substances have been utilised to treat infectious diseases including cardiac circulatory tonics and antiseptics as well as inflammation in humans.

(Mehta et al., 2003)

The plant's chief therapeutic characteristics include anticancer, ldl cholesterol reducing, antipyretic, antiinflammatory, antioxidant, antidiabetic, and fungicidal activities, among others. The flowers, leaves, bark, seeds, culmination, and roots are used to treat a variety of ailments.

(Gajjar et al., 2009)

When compared to the identical medicine nitrofurazone, embeliaribes showed a greater region of inhibition against the test organism Bacillus subtilis.

Aim and Objective

AIM- Aim of this Research work is to develop a herbal Nasal formulation

Objectives-

1. The worldwide health issue of AR is connected to life quality. Almost 40% of people worldwide have AR. AR, an allergic inflammation of the nasal airways, is characterized by nasal discharge, sneezing, itching, and congestion.
2. Antihistaminic medications, mast cell stabilizers, topical corticosteroids, and immunotherapy are currently used as therapies. All of these medications are ineffective and frequently cause a variety of unfavorable side effects, such as anxiety, exhaustion, headaches, sleepiness, and bleeding.
3. Due to nasal low residence duration, which necessitates repeated administration and resulting in overdose, existing formulations are unable to give relief, which causes a problem with rebound congestion.
4. The treatment of allergic illnesses has showed promise with a number of medicinal plants. In order to treat AR in ova challenged mice model, we created and developed polyherbal nasal formulations in the current work. With the aforementioned goals being achieved.

Plan of work

1. .Literature Review
2. Plants are chosen based on both historical and contemporary facts.
3. Identification of the plant and removal of plant matter.
4. To conduct pre-formulation research by choosing the excipients based on GRAS (generally regarded as safe).
5. Making nasal drop and nasal in situ gel compositions.
6. To carry out formulation optimization and evaluation.
7. Pharmacological formulation research using the AR mouse model and evaluation.

Expected outcomes

In present era AR is a type of fungal infection and a new outcome after immunosuppression. New drugs which are available in this time are highly costly which is out of reach of normal peoples. So we will try a creating a formulation that solve this problems. Herbal Nasal formulation is has low cost and effective and no side-effects comparison chemical based formulation. We will try to creating a formulation highly effective in nasal fungal infection and having long residence time in nasal.

References

- Abe, R., &Ohtani, K. (2013). An ethnobotanical study of medicinal plants and traditional therapies on Batan Island, the Philippines. Journal of ethnopharmacology, 145(2), 554-565..
- Ajuogu, P. K., Mgbere, O. O., Bila, D. S., & McFarlane, J. R. (2019). Hormonal changes, semen quality and variance in reproductive activity outcomes of post pubertal rabbits fed Moringaoleifera Lam. leaf powder. Journal of ethnopharmacology, 233, 80-86
- Al Suleimani, Y. M., & Walker, M. J. (2007). Allergic rhinitis and its pharmacology. Pharmacology & therapeutics, 114(3), 233-260.
- Altaş, S., Kızıl, G., Kızıl, M., Ketani, A., &Haris, P. I. (2011). Protective effect of Diyarbakır watermelon juice on carbon tetrachloride-induced toxicity in rats. Food and Chemical Toxicology, 49(9), 2433-2438.
- Alving K, Malinovschi A. 2010. Basic aspects of exhaled nitric oxide. Exhaled Biomarkers 49: 1–31.
- Amaglo, N. K., Bennett, R. N., Curto, R. B. L., Rosa, E. A., Turco, V. L., Giuffrida, A &Timpo, G. M. (2010). Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree Moringaoleifera L., grown in Ghana. Food Chemistry, 122(4), 1047-1054.
- Amit, A., Saxena, V. S., Pratibha, N., Bagchi, M., Bagchi, D., &Stohs, S. J. (2003). Safety of a novel botanical extract formula for ameliorating allergic rhinitis. Toxicology mechanisms and methods, 13(4), 253-261.
- Amresh, G., Singh, P. N., & Rao, C. V. (2008). Toxicological screening of traditional medicine Laghupatha (Cissampelospareira) in experimental animals. Journal of ethnopharmacology, 116(3), 454-460.
- Amresh, G., Singh, P. N., & Rao, C. V. (2008). Toxicological screening of traditional medicine Laghupatha (Cissampelospareira) in experimental animals. Journal of ethnopharmacology, 116(3), 454-460.
- Annapurna, D., &Rathore, T. S. (2010). Direct adventitious shoot induction and plant regeneration of EmbeliaribesBurm F. Plant Cell, Tissue and Organ Culture (PCTOC), 101(3), 269-277.

- Annapurna, D., &Rathore, T. S. (2010). Direct adventitious shoot induction and plant regeneration of EmbeliaribesBurm F. Plant Cell, Tissue and Organ Culture (PCTOC), 101(3), 269-277.
- Anwar, F., Latif, S., Ashraf, M., &Gilani, A. H. (2007). Moringaoleifera: a food plant with multiple medicinal uses. Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives, 21(1), 17-25
- . Anwar, F., Ashraf, M., Bhanger, M.I., (2005). Interprovenance variation in the composition of Moringaoleifera oilseeds from Pakistan. Journal of American Oil Chemist's Society 82, 45–51.
- Anon. (1966) The pharmacopoeia of India. Government of India Press, Nasik, p 795
- Arabshahi-Delouee, S., Aalami, M., Urooj, A., &Krishnakantha, T. P. (2009). Moringaoleifera leaves as an inhibitor of human platelet aggregation. Pharmaceutical biology, 47(8), 734-739.
- Asare, N., Instanes, C., Sandberg, W. J., Refsnes, M., Schwarze, P., Kruszewski, M., &Brunborg, G. (2012). Cytotoxic and genotoxic effects of silver nanoparticles in testicular cells. Toxicology, 291(1-3), 65-72.
- Aswar, U. M., Kandhare, A. D., Mohan, V., &Thakurdesai, P. A. (2015). Anti-allergic effect of intranasal administration of type-A procyanidin polyphenols based standardized extract of cinnamon bark in ovalbumin sensitized BALB/c mice. Phytotherapy research, 29(3), 423-433.
- Bachert, C., Krysko, O., Gevaert, P., Berings, M., Perez-Novo, C., & Van Crombruggen, K. (2015). Nasal Immunity, Rhinitis, and Rhinosinusitis. In Mucosal Immunology (pp. 1899-1921). Academic Press.
- Bahekar, P.C., Shah, J.H., Ayer, U.B., Mandhane, S.N., Thennati, R., (2008). Validation of guinea pig model of allergic rhinitis by oral and topical drugs. International Immunopharmacology 8, 1540–1551.
- Basu S, Bandyopadhyay AK (2010). Development and characterization of mucoadhesive in-situ nasal gel of midazolam prepared with Ficuscarica mucilage. AAPS PharmSciTech;11:1223-31.

- Belgamwar, V. S., Chauk, D. S., Mahajan, H. S., Jain, S. A., Gattani, S. G., &Surana, S. J. (2009). Formulation and evaluation of in situ gelling system of dimenhydrinate for nasal administration. Pharmaceutical development and technology, 14(3), 240-248.
- Berger WE. Allergic rhinitis in children: diagnosis and management strategies. Paediatr Drugs 2004;6:233–250. ¬ Bertram, U., &Bodmeier, R. (2012). Effect of polymer molecular weight and of polymer blends on the properties of rapidly gelling nasal inserts. Drug development and industrial pharmacy, 38(6), 659-669.
- Bennett, R.N., Mellon, F.A., Kroon, P.A., (2004). Screening crucifer seeds as sources of specific intact glucosinolates using ion-pair high-performance liquid chromatography negative ion electrospray mass spectrometry. J. Agric. Food Chem. 52, 428–438.
- Bhalerao AV, Lonkar SL, Deshkar SS, Shirolkar SV, Deshpande AD(2009). Nasal mucoadhesive in-situ gel of ondansetron hydrochloride. Indian J Pharm Sci;71:711–3.
- Bhandwalkar M, Amelia M (2013). Thermoreversible Nasal In situ Gel of Venlafaxine Hydrochloride: Formulation, Characterization, and Pharmacodynamic Evaluation. AAPS PharmSciTech;14(1): 101-10.